

Efficient Management of Water, Nutrient Resources

Assessing potential for drip irrigation and fertigation

Dr. Fred E. Below, Dr. Tryston Beyer, Juliann Seebauer and Dr. Ross Bender

The Fluid Journal • Official Journal of the Fluid Fertilizer Foundation • Winter 2018 • Vol. 25, No. 4, Issue #99

▼ DOWNLOAD

Summary: The industrywide initiative of doubling corn (*Zea mays L.*) grain yields by 2030 is required to feed a growing world population. The use of precision irrigation and fertilizer application technology may serve as a promising venue for producers to increase yields sustainably. Our findings highlight that adequate nutrient availability is needed upfront in corn and that delaying half of the fertilization until after the V5 stage by fertigation may lead to less nutrient loss, but may hinder maximum crop production.



To feed a growing human population on less land, greater yields are necessary. While the average U.S. corn yield is approximately 170 bu/A, greater yields are possible as shown by the 2016 National Corn Growers Contest winners, all exceeding 300 bu/A and almost a third of them exceeding 400 bu/A. The world record for corn is now well over 500 bushels per acre. In growing a 260 bu/A corn crop, our laboratory has determined that weather conditions account for over 27% of those bushels, while controllable crop management factors of nitrogen (N) fertilizer, hybrid, previous crop, plant population, tillage and plant growth regulators, each account for, on average, 26, 19, 10, 8, 6, and 4 percent of yield respectively (Agron J. 107:495). These yield estimates are based on prerequisites of drainage, pest and weed control, proper soil pH, and adequate P and K based on soil tests. Yield

increases in corn in the past 30 years in the U.S. have paralleled increases in planting density. However, this increased plant density leads to less root mass per plant and a greater need for more precise nutrient supply.

A series of studies conducted over the past three years have identified the fertility requirements for high-yielding corn (Agron. J. 105:161) and soybeans (Agron. J. 107:563; Table 1). Total nutrient requirements for soybeans are similar to those for corn, despite the misconception among farmers that nutrient management in soybeans is less critical because of N fixation as well as the notion that fertilizer supplied to a corn crop will also meet subsequent soybean fertility requirements. Nutrient harvest index values (i.e., the portion of total nutrient uptake represented in grain tissues) of N, P, sulfur (S) and zinc (Zn) in both corn and soybeans are generally between 60- 80%,

which suggests that soil test levels will quickly decline if provided inadequate crop nutrition. This may partially explain the decreases in soil concentrations of P, K, S, and Zn reported recently by IPNI (Better Crops 94(4):6). Current corn and soybean fertility recommendations are based on expected (corn) yield for N, and soil test levels for P and K. However, in the future, maximizing yields while also sustaining the productivity of soils requires a comprehensive season-long fertility plan designed to meet the uptake needs of well-managed corn and soybean crops.

Subsurface drip irrigation (SDI) has traditionally been used in the production of high value vegetable and fruit crops such as tomatoes. More recently, it has been adapted for commodity row crops such as corn, soybeans, and cotton as an alternative to center pivot or flood irrigation. The benefits of a SDI system relative to other traditional

irrigation forms include reduced water use brought about by up to a 50% reduction in evaporation losses (Irrig. Sci. 22:195L), and the ability to adapt to any field size, geometry or topography (Netafimusa.com). Additionally, SDI provides the opportunity to increase the efficiency of nutrient applications through the practice of fertigation (i.e., fluid fertilizer sources supplied with irrigation water). Fertigation of nutrients directly into the root microenvironment, particularly during periods of rapid uptake, can minimize nutrient losses associated with immobilization, volatilization, or surface runoff (Hortic. Technol. 6:(3):168). Currently, fertigated corn and soybean acreage in Illinois is limited in scope (90,000 acres or less than 1% of total crop acres as of a USDA-NASS 2013 survey). However, factors may accelerate fertigation and SDI system adoption in traditionally non-irrigated parts of the Corn Belt, including 1) high commodity and input prices, 2) catastrophic weather events such as the 2012 drought, and 3) the demand for increased agricultural productivity in response to world population growth.

Macronutrient accumulation varies considerably among crop and mineral element. In corn, the majority of N and K accumulation occurs before flowering compared to uptake of P, S, and Zn, which primarily occurs during grain filling (Agron. J 105:161). Because nutrient applications for corn production primarily supply nutrients in a bulk, dry granular form prior to planting, many nutrients are prone to chemical conversion or are fixed into unavailable forms before plant uptake. The potential for nutrient fixation or environmental pollution occurs for soybeans as well, with an estimated 118, 19, and 48 lbs/A, respectfully, of N P₂O₅ and K₂O uptake required after the initiation of pod-filling (Agron. J. 107:563).

In our initial studies using fertigation in 2014, we observed

Table 1. Mineral nutrition required to produce 230 bu acre⁻¹ corn (adapted from Bender et al., 2013) and 60 bu acre⁻¹ soybean (adapted from Bender et al., 2015). 'Maximum Uptake' (total nutrient uptake), 'Removal with Grain' (nutrient content of grain), and 'Harvest Index' (portion of total nutrient uptake residing in grain) are three key measures used to estimate nutritional needs in a cropping system.

Nutrient	Maximum Uptake		Removal With Grain		Harvest Index	
	Corn	Soybean	Corn	Soybean	Corn	Soybean
N	256	245	148	179	58	73
P ₂ O ₅	101	43	80	35	79	81
K ₂ O	180	170	59	70	32	41
S	23	17	13	10	57	61
Zn (oz ac ⁻¹)	7.1	4.8	4.4	2.0	62	44

Table 2. Nutrient application schedule for corn fertigation at Champaign, IL in 2016.

Growth Stage	Application Amount/ Cumulative Total†		
	N	P ₂ O ₅	K ₂ O
	----- lbs/ acre -----		
V5 - V6	45/ 45	0/ 0	0/ 0
V7 - V8	9/ 54	0/ 0	10/ 10
V11 - V12	9/ 63	10/ 10	15/ 25
V13 - V14	9/ 72	10/ 20	10/ 35
VT - R1	5/ 77	0/ 20	10/ 45
R2 - R3	5/ 82	10/ 30	5/ 50
R4 - R5	8/ 90	20/ 50	0/ 50

† In addition to 90 lbs N, 50 lbs P₂O₅, 50 lbs K₂O per acre at planting.

that even when water from irrigation is not needed, corn yields can be increased by 20 bushels per acre and soybean yields by 5 bushels per acre by better timing nutrient availability with plant needs. The importance of supplying nutrients at key growth stages may be more crucial for intensively managed corn and soybean production systems where other factors such as germ-plasm, pest control, plant density, and row spacing have been optimized. This project is designed to be forward-looking and assess how drip irrigation and fertigation might be used in the future to increase corn and soybean yields while also improving nutrient use efficiency.

Objectives

Our primary objectives were to 1) investigate subsurface drip irrigation (SDI) as a possible strategy to improve the efficiency of nutrient uptake and use when fluid nutrients are applied at key growth stages in corn and soybeans (*Glycine max* (L.) Merr.), and 2) understand how drip irrigation and fertigation can be optimized in a high yield agronomic

system with complementary agronomic management practices, including hybrid and variety selection and crop protection.

Methodology

Sites, cultural practices. Experiments were conducted at Champaign, IL using adjacent plots maintained in a corn/soybean rotation. The fertigation plots are situated on a Drummer-Flanagan Association soil (silt loam, 3.6% organic matter, 21.1 meq/100g CEC, 5.8 pH, 22 ppm P, 99 ppm K, and 9.5 ppm S using Mehlich 3 extraction) that is tile drained. The subsurface drip irrigation (SDI) fertigation system was installed in the spring of 2014 and consists of 48 equally sized zones that can be regulated for differential application of irrigation and fertigation. Using programmable controllers, each zone can precisely supply varying rates of nutrients at specific growth stages, according to plant needs. The dripper lines are buried approximately 14 to 16 inches below the soil surface with 30-inch spacing between lines. The SDI system covers ten acres total, divided

equally between corn and soybeans. Plots were maintained weed- and disease-free.

Research approach. Five corn hybrids (Croplan 8621 VT2P, Croplan 7087 VT2P, DeKalb C61-54, DeKalb C64-87, and NK 74R-3000GT) were planted on 21 May 2016. Using a precision plot planter (SeedPro 360, ALMACO, Nevada, IA) four population intensities were evaluated (32,000, 38,000, 44,000, and 50,000 plant/A). Con plots were four rows wide and 37.5 feet in length with 30-inch row spacing.

Nutrient applications. Half of the plots received a preplant broadcast application of 180 lbs N/A as urea, plus 100 lbs P₂O₅/A as Microessentials (MESZ, The Mosaic Company), plus 100 lbs K₂O/A as Aspire (The Mosaic Company) on May 2016. The remaining plots received half of the above rates as preplant, then equilibrated with the remaining 90 lbs N, 50 lbs P₂O₅, and 50 lbs K₂O/A during seven fertigation periods between V5 and R5 (Table 2). The values and the timing at which nutrients were acquired in a non-irrigated environment (Agron. J. 105:161), was the basis for the fertigation design (Table 2). Water was applied only as a medium for the fertigation component of this study. Soluble nutrient sources were used with a low ability to form precipitates in solution (e.g., urea ammonium nitrate, ammonium polyphosphate, ammonium thiosulfate, etc.). During periods of fertigation, a non-continuous (bulk) application of nutrients was applied during specific developmental stages (Table 2). An equal parts application technique was used in which 1) the system is brought to full pressure with water, 2) nutrients are injected and applied, and 3) the system is allowed to rinse with only irrigation water.

Measured parameters. Soil samples were obtained from plot areas prior to planting to confirm that fertility levels are uniform

across the site. Daily air and soil temperatures, precipitation, irrigation, and soil moisture were monitored throughout the growing season. Yield for corn was obtained on October 16, 2016 using a plot combine on the center two rows of each plot, and adjusted to 15.5% grain moisture concentration. Experimental units were arranged in a split-plot RCB design with six blocks. The main plot was designed with two fertilization levels: 100% preplant fertilizer and rain-fed only, versus 50% preplant fertilizer: 50% fertigated. The split plot was randomly assigned within each treatment block. Hybrid, fertilization/irrigation system, and population were considered fixed effects, while block and interactions with blocks were considered random effects. Measured parameters were analyzed using the PROC MIXED procedure of SAS (Version 8, SAS Institute, Cary, NC) and means were separated using Fisher's protected LSD test at the 0.10 level of significance.

Results

2014 and 2015. In 2014, due to a wet spring, the SDI system was completed behind schedule, and the corn crop wasn't planted until June 2014. Four hybrids were evaluated at populations ranging from 24,000 to 48,000 plants/A. A control treatment of 180 lbs N/A applied at V4 was compared to a fertigation system of an additional 80 lbs N, 0 lbs P₂O₅, 70 lbs K₂O, and 14 lbs S per acre applied between the V6 and R2 plant growth stages. With up to 44% of the fertigated nutrients recovered in the aboveground biomass, yields were increased about 10% over the irrigated control.

In 2015, more timely planting of five corn hybrids occurred on April 28, with populations ranging from 32,000 to 50,000 plants/A evaluated. An irrigated control received 180 lbs N/A at planting, although, due to abundant rainfall, only minimal irrigation was necessary during the season. The SDI system supplied

an additional 113 lbs N, 120 lbs P₂O₅, 150 lbs K₂O, 12 lbs S and 16 oz Zn per acre during seven fertigation periods between V5 to R5 in addition to a standard application of 180 lbs N/A as urea at planting. When averaged over the hybrids, fertigation increased yields over the irrigated control by 52 bu/A, achieving 246 bu/A. While there was a trend for greater corn yields with increasing population in the irrigated treatment, fertigation maximized yields even at the lowest planting population. All hybrids had increased yields in response to fertigation, varying from 49 to 56 bu/A.

2016. The initial studies in 2014 and 2015 provided a greater level of nutrients via fertigation, which resulted in greater yields. Therefore, in 2016, to investigate the potential for nutrient efficiency of fertigation, the objective was to provide the same total level of nutrients, 180 lbs N, 100 lbs P₂O₅, 100 lbs K₂O/A, to both fertigated and unfertigated corn, using fertigation to spread out the timing of nutrient availability commensurate with plant nutrient uptake. Compared to an all-upfront-preplant fertilized non-irrigated control, fertigated plots received half of the nutrient supply preplant, the other half between V5 And R5 (Table 2).

The 2016 production year experienced near-ideal growing conditions in Champaign, IL, with average temperatures and timely, adequate rainfall throughout the growing season (Table 3). As a result, corn in this region experienced little weather-induced heat or moisture stress and near record yields were chronicled for corn at the local state, regional, and/or national level. The fertigation system used the equivalent of 17 inches of water, similar to the May-August natural precipitation at Champaign, IL (Table 3).

When averaged over the five corn hybrids, the 100% preplant rainfed control produced yields of 247 bu/A

(Table 4). In contrast in 2016, by providing half fertilization preplant, a 50 percent delay in fertigation resulted in a 52 bu/A decrease in yield. Planting population did not change the overall results of either fertility program (Table 4). Notably, the 100% rainfed yields were similar to the fertigated yields of 2015, using less fertilizer.

Individual corn hybrid yields varied by 25 to 26 bu/A in both fertility scenarios but there were varietal differences (Table 5). The greatest yields in the 100% preplant control plots were produced by

7087VT2P and N74R-3000GT, while the greatest yields in the delayed fertigation were produced by DKC64-87 and 8621VT2PRIB (Table 5).

These yield decreases were due to the 10% decreases in both kernel number and kernel weight (daa not shown). Decreases in kernel number may be due to plant stresses before pollination. Potentially, the delay in N availability from fertilization in combination with the depth of the SDI system below the root zone, led to an early nutrient deficiency that was unrecoverable. However, these

Table 3. Temperature and precipitation during the production season at Champaign, IL in 2016 compared to the 30-year average. Data obtained from the Illinois State Water Survey.

Month	Precipitation (in)		Temperature (°F)	
	2016	30-Year Average	2016	30-Year Average
April	3.8	3.6	53	52
May	4.7	4.9	62	63
June	5.7	4.3	74	72
July	4.4	4.7	75	75
August	4.1	3.9	75	73
September	5.5	3.1	70	66

Table 4. Corn yield response to population, fertigation, and the difference of fertigation over 100% preplant fertilization at Champaign, IL in 2016. Values are averaged over five hybrids and six replications. All treatments received a total application of 180-100-100 lbs/acre of N- P₂O₅- and K₂O.

Population	100% preplant Rainfed	50% preplant: 50% Fertigated	Difference
----- bushels/ acre -----			
Plants/ acre			
32,000	246	195	-51*
38,000	249	195	-54*
44,000	244	195	-49*
50,000	250	195	-55*
Average	247	195	-52*

* Fertigated significantly different from 100% preplant at the 0.10 probability level.

Table 5. Corn yield response to hybrid, fertigation, and the difference of 50% fertigation over 100% preplant fertilization at Champaign, IL in 2016. Values are averaged over four populations and six replications. All treatments received a total application of 180-100-100 lbs/acre of N- P₂O₅- and K₂O.

Hybrid	100% preplant Rainfed	50% preplant: 50% Fertigated	Difference
----- bushels/ acre -----			
7087VT2P	258	197	-61*
8621VT2PRIB	244	200	-44*
DKC61-54	232	191	-41*
DKC64-87	248	205	-43*
N74R-3000GT	253	180	-73*
Average	247	195	-52*

* Fertigated significantly different from 100% preplant at the 0.10 probability level.

findings also indicate that some hybrids are more suited to fertigation and can better take advantage of the timed nutrient supply to produce greater yields with more efficient nutrient use.

Summing up

Using a subsurface drip irrigation system to precisely deliver supplemental fertilizer to the root zone successfully increased yields in both corn and soybeans in central IL up to by 52 bu/A for corn, depending on the hybrid. Supplying only half of the fertilizer preplant in 2016 led to yield losses, even when supplying the remainder throughout the growing season via fertigation. Typically, a greater planting population is necessary for increasing yields, however, the weather in all three years (2014 to 2016) enabled corn to maximize yield even at a population of 32,000 plants/A. Fertigation by SDI resulted in hybrid and variety yield differences, which suggests a need for characterization of optimal genetics and optimal nutrient supply scheduling for enhanced nutrient use and additional yield improvement.

Acknowledgements: The authors would like to acknowledge the generous support of numerous partners that helped make this research possible, including the Fluid Fertilizer Foundation, A & L Great Lakes Laboratories, AgVise Laboratories, BASF, John Deere, Mosaic, Monsanto, Netafim, Syngenta, USDA-NIFA project NC1200, Ward Laboratories, Inc., and Winfield United.

Dr. Below is Professor, Dr. Beyer is Research Specialist, and Mrs. Seebauer is Sr. Research Specialist in the Department of Crop Sciences, University of Illinois, West Gregory Drive, Urbana, IL 61802, and Dr. Bender is Sr. Agronomist, the Mosaic Company Cleveland, WI.